

Improved Algorithm for Bosonized Fermion Determinant.

A.A.Slavnov *

Steklov Mathematical Institute, Russian Academy of Sciences,
Vavilov st.42, GSP-1,117966, Moscow, Russia

February 1, 2008

Abstract

Following the line of ref. [1] we propose an improved algorithm which allows to calculate a D -dimensional fermion determinant integrating the exponent of $D + 1$ -dimensional Hermitean bosonic effective action. For a finite extra dimension the corrections decrease exponentially.

1 Introduction

Recently we proposed an algorithm which allows to calculate D -dimensional fermion determinants by integrating a $D + 1$ -dimensional bosonic effective action [1]. The following representation for the square of the covariant Dirac operator was obtained:

$$\begin{aligned} \det(\hat{D} + m)^2 = & \lim_{L \rightarrow \infty, b \rightarrow 0} \int \exp \{ a^4 b \sum_{n=-N+1}^N \sum_x [b^{-2}(\phi_{n+1}^*(x)\phi_n(x) + \phi_n^*(x)\phi_{n+1}(x) - 2\phi_n^*\phi_n) + \\ & + i[\phi_{n+1}^*(x)\gamma_5(\hat{D} + m)\phi_n(x) - \phi_n^*(x)\gamma_5(\hat{D} + m)\phi_{n+1}(x)][b]^{-1} + \\ & + \frac{1}{2}\phi_n^*(D^2 - m^2)\phi_n + \frac{i}{\sqrt{L}}(\phi_n^*(x)\chi(x) + \chi^*(x)\phi_n(x))] \} d\phi_n^* d\phi_n d\chi^* d\chi. \end{aligned} \quad (1)$$

Here ϕ_n are bosonic fields defined on a 4+1 dimensional lattice which have the same spinorial and internal structure as the Dirac fields ψ . The fifth component t to be defined on the one dimensional lattice of the length L with the lattice spacing b :

$$L = 2Nb, \quad -N < n \leq N \quad (2)$$

*E-mail: slavnov@class.mian.su

The free boundary conditions in t are assumed:

$$\phi_n = 0, \quad n \leq -N, \quad n > N \quad (3)$$

Lattice covariant derivative is denoted by D_μ

$$D_\mu \psi(x) = \frac{1}{a} [U_\mu(x) \psi(x + a_\mu) - \psi(x)] \quad (4)$$

$$\hat{D} = 1/2 \gamma_\mu (D_\mu^* - D_\mu) \quad (5)$$

(For simplicity we consider naive fermions, but all the construction is extended in a straightforward way to Wilson fermions).

For finite L and b the eq.(1) should be corrected by the terms

$$O(b^2 D^2) - \frac{4}{L^2 D^2} \sin^2\left(\frac{DL}{2}\right) \quad (6)$$

where D are eigenvalues of the Dirac operator.

One sees that although in the limit $b \rightarrow 0, L \rightarrow \infty$ the eq.(1) is exact, the convergence is not very fast. For numerical simulations it is important to make the corrections as small as possible, so one could use a one-dimensional lattice with relatively small number of sites N . Recently there were several publications, mainly based on the Lüscher's proposal [2] for bosonization of fermion determinants, where different aspects of numerical simulations in the bosonized models were discussed [3], [4] [5]. Further development in ref. [6], [7], [8] lead to algorithms with better convergence properties.

In the present paper I propose a modified algorithm, using essentially the same idea as in ref. [1], but providing much better convergence. Instead of polynomial suppression of finite size effects as in eq.(6), new algorithm provides exponential damping

$$\exp\{-mL\} \quad (7)$$

where m is a bare quark mass.

2 Improved algorithm for lattice QCD

Having in mind applications to QCD in this section we consider two fermion flavours interacting vectorially with the Yang-Mills fields U_μ . The reason to consider two degenerate flavours is the positivity of the square of the gauge covariant Dirac operator. It is convenient to present the square of fermion determinant in the following form

$$\begin{aligned} \int \exp\{a^4 \sum_{n=1}^2 \sum_x \bar{\psi}_n(x) (\hat{D} + m) \psi_n(x)\} d\bar{\psi} d\psi &= \det[\gamma_5 (\hat{D} + m)]^2 = \\ &= \int \exp\{a^4 \sum_x \bar{\psi}(x) (\hat{D}^2 - m^2) \psi(x)\} d\bar{\psi} d\psi \end{aligned} \quad (8)$$

We again introduce five dimensional bosonic fields $\phi(x, t)$ with the same spinorial and internal structure as $\psi(x)$. The notations are as above.

We are going to prove the following identity

$$\begin{aligned}
& \int \exp\{a^4 \sum_x \bar{\psi}(x)(\hat{D}^2 - m^2)\psi(x)\} d\bar{\psi} d\psi = \\
& = \lim_{L \rightarrow \infty, b \rightarrow 0} \int \exp\{a^4 b \sum_{n=-N+1}^N \sum_x [(b^{-2})(\phi_{n+1}^*(x)\phi_n(x) + h.c. - 2\phi_n^*\phi_n) + \\
& \quad -i\phi_{n+1}^*(x)\gamma_5 \hat{D} L^{-1}(2n+1)\phi_n + h.c. \\
& \quad + 1/2\phi_n^*(x)\hat{D}^2 L^{-2}(2n+1)^2 b^2 \phi_n \\
& \quad + i(\frac{2m}{\pi L^5})^{1/4}(\phi_n^*(x)\chi(x) + \chi^*(x)\phi_n(x))2nb \exp\{-mL^{-1}b^2 n^2\}]\} d\phi_n^* d\phi_n d\chi^* d\chi.
\end{aligned} \tag{9}$$

Integration goes over the fields $\phi_n(x)$ satisfying the free boundary conditions (3). The last term in the exponent in the r.h.s. of eq.(9) introduces the constraint

$$\sum_n \phi_n n \exp\{-mL^{-1}b^2 n^2\} = 0 \tag{10}$$

Substituting the solution of this constraint to the eq.(9) one gets the representation for the determinant of the gauge covariant Dirac operator as the path integral of the exponent of the bosonic Hermitean action. Below we present a construction which justifies the eq.(9) and estimate the corrections due to the finite lattice spacing b and finite lattice size L .

Our starting point is the integral

$$\begin{aligned}
I = \int \exp\{a^4 b \sum_{n=-N+1}^N \sum_x [(b^{-2})(\phi_{n+1}^*(x) \exp\{-i\gamma_5 \hat{D} b^2(n+1)^2 L^{-1}\} \times \\
\exp\{i\gamma_5 \hat{D} b^2 n^2 L^{-1}\} \phi_n(x) + h.c. - 2\phi_n^* \phi_n) + \\
+ i(\frac{2m}{\pi L^5})^{1/4}(\phi_n^*(x)\chi(x) + \chi^*(x)\phi_n(x))2nb \exp\{-mL^{-1}b^2 n^2\}]\} d\phi_n^* d\phi_n d\chi^* d\chi.
\end{aligned} \tag{11}$$

The operator $\gamma_5 \hat{D}$ is Hermitean and it's eigenvalues are real. R.h.s. of eq. (11) can be written in terms of eigenvectors of the operator $\gamma_5 \hat{D}$, which are denoted as ϕ_α :

$$\begin{aligned}
I = \int \exp\{b \sum_{n=-N+1}^N \sum_\alpha [(b^{-2})(\phi_{n+1}^{\alpha*} \exp\{-iD^\alpha b^2(n+1)^2 L^{-1}\} \times \\
\exp\{iD^\alpha b^2 n^2 L^{-1}\} \phi_n^\alpha + h.c. - 2\phi_n^{\alpha*} \phi_n^\alpha) + \\
+ i(\frac{2m}{\pi L^5})^{1/4}(\phi_n^{\alpha*} \chi^\alpha + \chi^{\alpha*} \phi_n^\alpha)2nb \exp\{-mL^{-1}b^2 n^2\}]\} d\phi_n^{\alpha*} d\phi_n^\alpha d\chi^{\alpha*} d\chi^\alpha
\end{aligned} \tag{12}$$

To calculate the integral (12) we make the following change of variables:

$$\phi_n^\alpha \rightarrow \exp\{-iD^\alpha n^2 b^2 L^{-1}\} \phi_n^\alpha, \quad \phi_n^{\alpha*} \rightarrow \exp\{iD^\alpha n^2 b^2 L^{-1}\} \phi_n^{\alpha*} \tag{13}$$

Then the integral (12) acquires the form

$$I = \int \exp\{b \sum_{n=-N}^N \sum_{\alpha} b^{-2} [\phi_{n+1}^{*\alpha} \phi_n^{\alpha} + \phi_n^{*\alpha} \phi_{n+1}^{\alpha} - 2\phi_n^{*\alpha} \phi_n^{\alpha}] +$$

$$+ i(\frac{2m}{\pi L^5})^{1/4} (\phi_n^{\alpha*} \exp\{iD^{\alpha} b^2 n^2 L^{-1}\} \chi^{\alpha} + \chi^{\alpha*} \exp\{-iD^{\alpha} b^2 n^2 L^{-1}\} \phi_n^{\alpha}) \times$$

$$2nb \exp\{-mL^{-1} b^2 n^2\} \} d\phi_n^{\alpha*} d\phi_n^{\alpha} d\chi^{\alpha*} d\chi^{\alpha}$$
(14)

Now the quadratic form in the exponent does not depend on D^{α} and therefore the corresponding determinant is a trivial constant. So we can calculate the integral by finding the stationary point of the exponent, which is defined by the following equations:

$$b^{-2}(\phi_{n+1}^{*\alpha} + \phi_{n-1}^{*\alpha} - 2\phi_n^{*\alpha}) +$$

$$i(\frac{2m}{\pi L^5})^{1/4} \chi^{*\alpha} 2nb \exp\{-(iD^{\alpha} + m)b^2 n^2 L^{-1}\} = 0, \quad n \neq -N$$

$$b^{-2}(\phi_{n+1}^{\alpha} + \phi_{n-1}^{\alpha} - 2\phi_n^{\alpha}) +$$

$$+ i(\frac{2m}{\pi L^5})^{1/4} \chi^{\alpha} 2nb \exp\{(iD^{\alpha} - m)b^2 n^2 L^{-1}\} = 0, \quad n \neq -N$$

$$b^{-2}(\phi_{-N+1}^{*\alpha} - 2\phi_{-N}^{*\alpha}) + i(\frac{2m}{\pi L^5})^{1/4} \chi^{*\alpha} 2Nb \exp\{-(iD^{\alpha} + m)b^2 N^2 L^{-1}\} = 0$$

$$b^{-2}(\phi_{-N+1}^{\alpha} - 2\phi_{-N}^{\alpha}) + i(\frac{2m}{\pi L^5})^{1/4} \chi^{\alpha} 2Nb \exp\{(iD^{\alpha} - m)b^2 N^2 L^{-1}\} = 0$$

$$\phi_{-N}^{\alpha} = \phi_{-N}^{\alpha*} = 0, \quad \phi_{N+1}^{\alpha} = \phi_{N+1}^{\alpha*} = 0$$
(15)

These equations are most easily solved for small b , when they can be approximated by the differential equations:

$$\ddot{\phi}^{*\alpha} + i(\frac{2m}{\pi L^5})^{1/4} \chi^{*\alpha} 2t \exp\{-(iD^{\alpha} + m)t^2 L^{-1}\} = 0$$

$$\ddot{\phi}^{\alpha} + i(\frac{2m}{\pi L^5})^{1/4} \chi^{\alpha} 2t \exp\{(iD^{\alpha} - m)t^2 L^{-1}\} = 0$$

$$\phi^{\alpha}(\frac{L}{2}) = \phi^{\alpha}(-\frac{L}{2}) = 0, \quad \phi^{\alpha*}(\frac{L}{2}) = \phi^{\alpha*}(-\frac{L}{2}) = 0$$
(16)

The solution of these eq.s is

$$\phi_{st}^{*\alpha} = i\chi^{*\alpha} \frac{(2m\pi L)^{1/4}}{2(m + iD^{\alpha})^{3/2}} \Phi(tL^{-1/2}(m + iD^{\alpha})^{1/2}) +$$
(17)

$$+ i\chi^{\alpha*} \frac{(2m)^{1/4}}{(\pi L)^{1/4}(m + iD^{\alpha})^2} \exp\{\frac{-L(m + iD^{\alpha})}{4}\} 2tL^{-1} - i\chi^{\alpha*} \frac{(2\pi mL)^{1/4}}{2(m + iD^{\alpha})^{3/2}}$$

$$\phi_{st}^{\alpha} = i\chi^{\alpha} \frac{(2m\pi L)^{1/4}}{2(m - iD^{\alpha})^{3/2}} \Phi(tL^{-1/2}(m - iD^{\alpha})^{1/2}) +$$
(18)

$$+i\chi^\alpha \frac{(2m)^{1/4}}{(\pi L)^{1/4}(m-iD^\alpha)^2} \exp\left\{\frac{-L(m-iD^\alpha)}{4}\right\} 2tL^{-1} - i\chi^\alpha \frac{(2\pi mL)^{1/4}}{2(m-iD^\alpha)^{3/2}}$$

where $\Phi(x)$ is a Fresnel integral. Substituting this solution into the integral (15) we get

$$I = \int \exp\left\{-\sum_\alpha \int_{-L/2}^{+L/2} dt \phi_{st}^{\alpha*}(t) \ddot{\phi}_{st}^\alpha(t)\right\} d\chi^{\alpha*} d\chi^\alpha \quad (19)$$

Integrating by parts and using the fact that $\phi_{st}^\alpha(L/2) = \phi_{st}^\alpha(-L/2) = 0$, we can transform this integral to the form

$$\begin{aligned} I = \int \exp\left\{\sum_\alpha \int_{-L/2}^{L/2} dt \dot{\phi}_{st}^{\alpha*} \dot{\phi}_{st}^\alpha\right\} d\chi^{\alpha*} d\chi^\alpha = \\ \int \exp\left\{-\sum_\alpha \left(\frac{2m}{\pi L}\right)^{1/2} \frac{\chi^{\alpha*} \chi^\alpha}{m^2 + (D^\alpha)^2} \left[\int_{-L/2}^{L/2} dt \exp\{-2mL^{-1}t^2\} + \right. \right. \\ \left. \left. + \frac{4}{L(m^2 + (D^\alpha)^2)} e^{\{-\frac{Lm}{2}\}} + \frac{2 \exp\{-\frac{L(m+iD^\alpha)}{4}\}}{(m+iD^\alpha)L} \int_{-L/2}^{L/2} dt \exp\{-(m-iD^\alpha)L^{-1}t^2\} + \right. \right. \\ \left. \left. \frac{2 \exp\{-\frac{L(m-iD^\alpha)}{4}\}}{(m-iD^\alpha)L} \int_{-L/2}^{L/2} dt \exp\{-(m+iD^\alpha)L^{-1}t^2\}\right\}\right\} \end{aligned} \quad (20)$$

Integrating over χ one gets

$$I = \prod_\alpha (m^2 + (D^\alpha)^2) (1 + O((mL)^{-3/2} \exp\{-\frac{mL}{2}\})) \quad (21)$$

So if $m \sim a^{-1}$, and $L = 2Nb$, the corrections are of the order

$$O((Nba^{-1})^{-3/2} \exp\{-Nba^{-1}\}) \quad (22)$$

At the same time it is easy to show that replacing the sum in the eq.(15) by the integral over t produces corrections of order $O(b^2a^{-2})$. Therefore taking $N \sim 3ab^{-1}$ we can make the corrections due to the finite size of a lattice less than 1 %.

Taking into account that

$$(\gamma_5 \hat{D})^2 = -(\hat{D})^2 \quad (23)$$

we see that

$$I = \det(\hat{D} + m)^2 + O(b^2a^{-2}) + O((mL)^{-3/2} \exp\{-\frac{mL}{2}\}) \quad (24)$$

To conclude we note that our equation (9) is a linearized version of the integral (11). The exponent in the quadratic form of the action (11) may be written in the form

$$\exp\{-i\gamma_5 \hat{D} b^2 L^{-1} (2n+1)\}, \quad 2nb \leq L \quad (25)$$

Having in mind that

$$\|\gamma_5 \hat{D}\| \leq 8a^{-1} \quad (26)$$

one sees that

$$\frac{\|\gamma_5 \hat{D} b^2 (2n+1)\|}{L} \leq 8ba^{-1} \quad (27)$$

and if $8b < a$ one can replace the exponential in the action (11) by the first terms of it's Taylor series, leading to the eq.(9).

3 Discussion

We proved that the purely bosonic integral (9) is equal to the square of the determinant of covariant Dirac operator up to the terms

$$O(b^2 a^{-2}) + O((mL)^{-\frac{3}{2}} \exp\{-\frac{mL}{2}\}) \quad (28)$$

Convergence of this approximation is much faster than in the ref. [1], where the finite size effects were suppressed polynomially. For finite lattice spacing b the corrections decrease exponentially when the number of extra fields N increases. To get the same accuracy one needs much smaller number of bosonic fields, which hopefully will simplify Monte-Carlo simulations. It is worthwhile to note that to get exponential damping it is crucial to introduce the exponential factor into the constraint (10). As for the choice of the quadratic form in the effective bosonic action, there exists a certain freedom. We choose the particular form (11) to simplify analytic calculations. Presumably the choice of the quadratic form done in our previous paper [1] is also possible. We also note that contrary to our previous algorithm [1] the present construction cannot be reduced to Lüscher's algorithm with a particular choice of a polynomial due to explicit dependence of the constraint on the mass.

Acknowledgements.

The main part of this work was done while the author was visiting Max-Planck Institute for Physics in Munich and Universite Paris-Sud Centre d'Orsay. I am grateful to the Theoretical Departments of these Institutes, and in particular to D.Maison and Ph.Boucaud for hospitality and interesting discussions. I thank A.Galli for helpful comments. This research was supported in part by Russian Basic Research Fund under grant 96-01-00551.

References

- [1] A.A.Slavnov SMI-28-95 hep-th/ 9512101,
- [2] M.Lüscher Nucl.Phys. B418 (1994) 637.
- [3] B.Bunk, K.Jansen, B.Jegerlehner, M.Lüscher, H.Simma, R.Sommer, Nucl.Phys.B (Proc.Suppl.)42 (1995) 49,
- [4] C.Alexandrou, A.Borelli, Ph. de Forcrand, A.Galli and F.Jegerlehner, Nucl.Phys B456 (1995) 296,
- [5] K.Jansen, B.Jegerlehner, C.Liu hep-lat/9604016,
- [6] I.Montvay, DESY 95-192, hep-lat/9510042,
- [7] A.Borici, Ph.de Forcrand IPS-95-23, hep-lat/9509080,
- [8] A.Borelli, Ph.de Forcrand, A.Galli, hep-lat/9602016